

Diagnostic tools for approximate Bayesian computation using the coverage property

D. Prangle*, M. G. B. Blum[†], G. Popovic[‡] and S. A. Sisson^{‡§}

Abstract

Approximate Bayesian computation (ABC) is an approach for sampling from an approximate posterior distribution in the presence of a computationally intractable likelihood function. A common implementation is based on simulating model, parameter and dataset triples, (m, θ, y) , from the prior, and then accepting as samples from the approximate posterior, those pairs (m, θ) for which y , or a summary of y , is “close” to the observed data. Closeness is typically determined through a distance measure and a kernel scale parameter, ϵ . Appropriate choice of ϵ is important to producing a good quality approximation. This paper proposes diagnostic tools for the choice of ϵ based on assessing the coverage property, which asserts that credible intervals have the correct coverage levels. We provide theoretical results on coverage for both model and parameter inference, and adapt these into diagnostics for the ABC context. We re-analyse a study on human demographic history to determine whether the adopted posterior approximation was appropriate. R code implementing the proposed methodology is freely available in the package `abc`.

Keywords: Approximate Bayesian computation; Coverage; Diagnostics; Model inference; Likelihood-free inference.

*Mathematics and Statistics Department, Lancaster University, Lancaster, U.K.

[†]Université Joseph Fourier, Centre National de la Recherche Scientifique, Laboratoire TIMC-IMAG UMR 5525, Grenoble, F-38041, France

[‡]School of Mathematics and Statistics, University of New South Wales, Sydney, Australia

[§]Email: `Scott.Sisson@unsw.edu.au`

1 Introduction

For a given model, m , Bayesian inference for unknown parameters θ , and given observed data y_{obs} , updates prior beliefs $\pi(\theta|m)$ through the likelihood function $\pi(y_{\text{obs}}|\theta, m)$. This produces the posterior distribution $\pi(\theta|y_{\text{obs}}, m) = \pi(y_{\text{obs}}|\theta, m)\pi(\theta|m)/\pi(y_{\text{obs}}|m)$, where $\pi(y_{\text{obs}}|m) = \int \pi(y_{\text{obs}}|\theta, m)\pi(\theta|m)d\theta$ is the integrated likelihood for model m . Similarly, Bayesian inference for a discrete set of models $m \in \{1, 2, \dots, M\}$ updates a prior mass function $p(m)$ to posterior weights $p(m|y_{\text{obs}}) \propto \pi(y_{\text{obs}}|m)p(m)$. Here the joint posterior of parameter and model is given by $\pi(\theta, m|y_{\text{obs}}) \propto \pi(y_{\text{obs}}|\theta, m)\pi(\theta|m)p(m)$. Increased usage of Bayesian inference in recent decades has been built on powerful algorithms, such as Markov chain Monte Carlo, which make use of repeated evaluation of the likelihood function(s).

Approximate Bayesian computation (ABC) refers to a family of algorithms which perform an approximate Bayesian inference when numerical evaluation of the likelihood is not feasible, but where it is possible to simulate from the model(s) $y \sim \pi(\cdot|\theta, m)$. ABC has become a popular tool for the analysis of complicated models in a wide range of challenging applications. See e.g. Beaumont (2010); Bertorelle et al. (2010); Csilléry et al. (2010); Marin et al. (2012) and Sisson and Fan (2011) for overviews of methods and applications.

A common, importance sampling-based implementation of ABC, expressed for the multi-model setting, is given by the following:

ABC importance sampling

1. For $i = 1, \dots, N$:
 Sample a model and parameter from the prior $(\theta_i, m_i) \sim \pi(\theta|m)p(m)$.
 Simulate data from model m_i as $y_i \sim \pi(y|\theta_i, m_i)$.
 2. Weight each sample (y_i, θ_i, m_i) by $w_i \propto K_\epsilon(\|y_i - y_{\text{obs}}\|)$.
-

Algorithm 1: An ABC importance sampling algorithm, based on a single large sample of size N , incorporating model choice and parameter inference.

Here $K_\epsilon(u) = K(u/\epsilon)/\epsilon$ is a standard smoothing kernel with scale parameter $\epsilon > 0$, and $\|\cdot\|$ is a distance measure e.g. Euclidean. For this article, for simplicity and w.l.o.g.

we consider $K_\epsilon(u)$ to be the uniform kernel $U(-\epsilon, \epsilon)$, so that step 2 above corresponds to selecting (i.e. with non-zero weights) those samples (y_i, θ_i, m_i) for which $\|y_i - y_{\text{obs}}\| \leq \epsilon$. Note that when $p(m)$ is small, the above algorithm may have large Monte Carlo error in estimating $p(m|y_{\text{obs}})$. This is often avoided by using a uniform prior mass function in place of $p(m)$ as a computational device, and then reweighting each model appropriately (see e.g. Grelaud et al. (2009)).

The output of the above algorithm is a sample of parameter vectors (θ_i, m_i) from an approximation to the posterior

$$\pi_{\text{ABC}}(\theta, m|y_{\text{obs}}) \propto \pi(\theta|m)p(m) \int \pi(y|\theta, m) K_\epsilon(\|y - y_{\text{obs}}\|) dy, \quad (1)$$

where it can be seen that $\pi_{\text{ABC}}(\theta, m|y_{\text{obs}}) \approx \pi(\theta, m|y_{\text{obs}})$ following standard conditional density estimation arguments. There are two sources of approximation error in the sample representation of (1) from Algorithm 1, both of which are influenced by ϵ . The first is the discrepancy between $\pi_{\text{ABC}}(\theta, m|y_{\text{obs}})$ and $\pi(\theta, m|y_{\text{obs}})$. These are equal in the limit $\epsilon \rightarrow 0$, however the approximation deteriorates as ϵ is increased. The extreme result is $\lim_{\epsilon \rightarrow \infty} \pi_{\text{ABC}}(\theta, m|y_{\text{obs}}) = \pi(\theta|m)p(m)$. That is, all proposed samples (θ, m) are accepted, so that ABC targets the prior distribution and all information from the data is lost. Secondly, $\pi_{\text{ABC}}(\theta, m|y_{\text{obs}})$ is approximated by a finite sample whose size reduces as $\epsilon \rightarrow 0$ (in other ABC algorithms this corresponds to extremely low acceptance rates). Indeed the sample size typically reduces to zero for continuous data. In effect, ϵ controls a form of the usual bias-variance trade-off (Blum, 2010a).

Many approaches have been proposed to reduce the approximation error. One is to replace $\|y - y_{\text{obs}}\|$ by $\|s - s_{\text{obs}}\|$ where $s = S(y)$ is a vector of summary statistics. Low dimensional but informative summary statistics can greatly improve inferential accuracy, even at the price of potential information loss (Blum et al., 2013). Using summary statistics introduces another level of approximation, as then ABC approximates $\pi(\theta, m|s_{\text{obs}})$ rather

than $\pi(\theta, m|y_{\text{obs}})$. A second approach is to post-process the sample from $\pi_{\text{ABC}}(\theta, m|y_{\text{obs}})$ with $\epsilon > 0$ so that is approximately transformed to a sample with $\epsilon = 0$. Termed regression-adjustment, for within-model parameter inference this takes the form of linear or non-linear regression-based transformations of θ_i (Beaumont et al., 2002; Blum and François, 2010; Blum et al., 2013). For the adjustment of model probabilities, post-processing can be performed by multinomial regression (Beaumont, 2008).

1.1 Diagnostics for ABC

This paper addresses two related open questions about ABC in practice. Firstly, is it possible to validate the ABC approximation of the posterior, $\pi_{\text{ABC}}(\theta, m|y_{\text{obs}}) \approx \pi(\theta, m|y_{\text{obs}})$ (or $\pi_{\text{ABC}}(\theta, m|s_{\text{obs}}) \approx \pi(\theta, m|s_{\text{obs}})$ when using summary statistics), as accurate? Secondly, how should ϵ be chosen? Typically ϵ is commonly chosen in an ad-hoc manner, although several authors (Bortot et al., 2007; Ratmann et al., 2009; Blum, 2010b; Faisai et al., 2013) have suggested approaches where ϵ is estimated as part of an extended model.

In this paper we approach the question of the accuracy of the ABC posterior approximation by examining whether the *coverage property* holds (described below). By numerically evaluating adherence to the coverage property through diagnostic statistics, we are able to determine the likely accuracy of the ABC approximation for a range of ϵ values. In favourable circumstances this allows the user to choose as large a value of ϵ as possible (for computational efficiency) for which coverage approximately holds. Alternatively, the coverage diagnostics may reveal that large approximation error remains for any choice of ϵ .

Our approach is based on credible intervals, a standard Bayesian method to give interval parameter estimates. Consider the case of within-model parameter inference for a fixed model m_0 . An $\alpha\%$ credible interval for a univariate parameter θ is an interval I with the property that $\Pr(\theta \in I|y_{\text{obs}}) = \alpha/100$. Suppose that credible intervals are constructed from ABC output for data simulated from a known parameter value, θ_0 . Roughly speaking, the coverage property asserts that these intervals have the claimed probability of containing θ_0 .

To fully define the property we must be specific about the distribution of θ_0 . This is discussed later, where we give theoretical results supporting a particular choice for our purpose. We also describe how this definition can be extended to model inference problems.

An equivalent condition to the coverage property was provided by Cook et al. (2006). Namely, the p -values of $\theta_0 \leq \theta$ (or $\theta_0 > \theta$) within the posterior estimates must have a $U(0, 1)$ distribution. Accordingly, the coverage property may be tested for a particular value of ϵ , by repeatedly performing ABC for many choices of θ_0 and associated pseudo-observed data $y_0 \sim \pi(y|\theta_0, m_0)$, computing p -values, and then applying standard tests for uniformity. We provide a similar equivalent condition and a test of the coverage property for model probabilities, in addition to providing a computationally efficient process to compute the test statistics, for both within- and between-models.

There is a sizeable literature related to the coverage property. Bayesian work includes determining the correctness of complex Bayesian simulation algorithms (Cook et al., 2006), the post-processing of within-model ABC output (Menéndez et al., 2012) and the validation of ABC analyses in the single model setting (Wegmann et al., 2009, 2010; Aeschbacher et al., 2012). A recent overview of work from a frequentist perspective is provided by Gneiting et al. (2007). However, this work has the somewhat different aim of determining consistency between statistical predictions and a sequence of observed outcomes (e.g. weather forecasts and meteorological data). Despite the difference in aims, the primary ideas behind our statistical tests of coverage go back to the frequentist literature: Dawid (1984) for continuous parameters and Seillier-Moiseiwitsch and Dawid (1993) for model choice. The approach we develop in this article is similar to the ABC papers mentioned above. Our contribution here, is to explain and justify the theoretical basis and methodology of coverage in more detail, to improve this methodology where needed, and to extend these ideas to the hitherto unconsidered realm of model inference for ABC.

The remainder of the paper is structured as follows: Section 2 defines the coverage property for both parameter and model inference, and gives some theoretical results. The

methodological details of the resulting diagnostics are described in Section 3, including an algorithm and discussion of diagnostic statistics and tests. Section 4 presents a simulated example to illustrate the methods and justify some implementation choices, followed by a re-analysis of a study into human demographic history to determine whether a reliable posterior approximation was obtained. Finally, Section 5 concludes with a discussion.

2 Coverage

We investigate whether the ABC approximation $\pi_{\text{ABC}}(\theta, m|y_{\text{obs}})$ (or $\pi_{\text{ABC}}(\theta, m|s_{\text{obs}})$) is a good representation of the posterior $\pi(\theta, m|y_{\text{obs}})$ (or $\pi(\theta, m|s_{\text{obs}})$) by testing the *coverage property*. For inference on a continuous scalar parameter, θ , an informal definition is that a given credible interval based on $\theta|y_0$, where $y_0 \sim \pi(y|\theta_0, m_0)$ for fixed m_0 , should contain the true parameter, θ_0 , the appropriate proportion of times. This Section presents a more precise definition, a discussion of the property’s consequences, and results on how it can be tested. We also describe a version of the property suitable for a model choice setting. We notationally work with y rather than s throughout this Section.

2.1 Parameter inference

We define the coverage property for the case of a continuous scalar parameter θ for a fixed model m_0 , where for the remainder of Section 2.1 we drop all notational dependence on m_0 . For multivariate parameter vectors, our method will examine each parameter separately.

The informal definition above is based on analysing data y_0 simulated from known parameter values θ_0 . To formalise the property we introduce a distribution for these, $H(\theta_0, y_0)$, with densities associated with H denoted by h . A natural choice, used by Cook et al. (2006), Wegmann et al. (2009, 2010) and Aeschbacher et al. (2012), is $h(\theta_0, y_0) = \pi(y_0|\theta_0)\pi(\theta_0)$; draw the parameters from the prior, and the data from the model of interest conditional on this. We present an argument in favour of an alternative choice for the ABC setting below.

Let $g(\theta|y)$ be a density approximating the posterior given data y . From this, credible intervals of any level can be constructed. Suppose $C(y, \alpha)$ is a credible interval of level $\alpha\%$. We say that g satisfies coverage with respect to H if the coverage level of such an interval is α when analysing data generated from H (i.e. $\Pr(\theta_0 \in C(y_0, \alpha)) = \alpha$), for any choice of α and C . More formally, we have:

Definition of coverage property: Let $g(\theta|y)$ be a density approximating the univariate posterior $\pi(\theta|y)$, and $G_y(\theta)$ be the corresponding distribution function. Consider a function $B(\alpha) \subseteq [0, 1]$ defined for $\alpha \in [0, 1]$ such that the resulting set has Lebesgue measure α . Let $C(y, \alpha) = G_y^{-1}[B(\alpha)]$ and $H(\theta_0, y_0)$ be the distribution function for (θ_0, y_0) . We say g satisfies the coverage property with respect to distribution $H(\theta_0, y_0)$ if for every function B and every $\alpha \in [0, 1]$, $\Pr(\theta_0 \in C(y_0, \alpha)) = \alpha$.

There are two requirements for the coverage property to be a useful criterion to determine how well $g(\theta|y)$ approximates the posterior $\pi(\theta|y)$. These requirements will determine some characteristics of $H(\theta_0, y_0)$. Firstly, it should hold when $g(\theta|y) = \pi(\theta|y)$.

Result 1 *The posterior, $\pi(\theta|y)$, satisfies coverage with respect to any distribution $H(\theta_0, y_0)$ with conditional density $h(\theta_0|y_0) = \pi(\theta_0|y_0)$.* Proof in Appendix.

The second requirement is that the coverage property should avoid false positives: it should not hold when $g(\theta|y) \neq \pi(\theta|y)$. However, coverage can hold when $g(\theta|y) = \pi(\theta)$ equals the prior distribution, when $\theta_0 \sim \pi(\theta)$ is also drawn from the prior.

Result 2 *The prior, $\pi(\theta)$, satisfies coverage with respect to any distribution $H(\theta_0, y_0)$ with marginal density $h(\theta_0) = \pi(\theta_0)$.* Proof in Appendix.

The above results demonstrate that the choice $h(\theta_0, y_0) = \pi(\theta_0, y_0)$ (where $\pi(\theta_0, y_0) = \pi(\theta_0|y_0)\pi(y_0) = \pi(y_0|\theta_0)\pi(\theta_0)$) leads to the coverage property holding for both the prior and posterior distributions. The false positive of the prior is particularly unwelcome in the ABC

context, as it corresponds exactly to the ABC approximation $\pi_{\text{ABC}}(\theta|y)$ for $\epsilon \rightarrow \infty$. (The prior also coincides with the ABC posterior approximation when $s = S(y)$ has no information for θ under the model, so that $\pi_{\text{ABC}}(\theta|s) = \pi(\theta|s) = \pi(\theta)$, although this is due to Result 1 rather than Result 2. See Section 5 for more discussion.)

To avoid this we propose the alternative choice of $h(\theta_0, y_0) \propto \pi(\theta_0, y_0)\mathbb{I}[y_0 \in A]$. That is, the distribution $\pi(\theta_0, y_0)$ truncated to require that the data lie within some subset A . This preserves $h(\theta_0|y_0) = \pi(\theta_0|y_0)$, so coverage holds for the posterior, but it typically alters $\pi(\theta_0)$ so that coverage no longer holds for the prior (i.e. $h(\theta_0) \neq \pi(\theta_0)$). We examine some convenient choices of A in Section 3. In this manner, we are aiming to evaluate coverage for datasets similar to y_{obs} , rather than the much stronger context of coverage holding for all datasets. Note that the above results do not prove that the posterior $\pi(\theta|y)$ is the only distribution to satisfy coverage with respect to our choice of H . However, we are unaware of any other such distributions that are likely to arise in the ABC context.

An equivalent condition to the coverage property, which is easier to test, is the following:

Result 3 *Let H be the distribution function of (θ_0, y_0) . Define $p_0 = G_{y_0}(\theta_0)$, where $G_y(\theta)$ is the distribution function of θ under $g(\theta|y)$. Coverage holds with respect to H iff*

$$p_0 \sim U(0, 1). \tag{2}$$

Proof in Appendix.

A similar result was given by Cook et al. (2006), who proved that under coverage (with respect to the distribution function $h(\theta_0, y_0) = \pi(y_0|\theta_0)\pi(\theta_0)$) the empirical distribution of p_0 converges to $U(0, 1)$.

2.2 Model inference

As for parameter inference, the definition of the coverage property for model inference requires us to specify the distribution of the known parameter values m_0 and y_0 through

$H(m_0, y_0)$, which can be considered a marginal distribution of $H(m_0, \theta_0, y_0)$. Note that for model inference, H , its derivatives (h), and the mass function $g(m|y)$ approximating the posterior $p(m|y)$ are discrete functions.

Formalising the intuitive notion of coverage for the case of model choice faces the difficulty of interpreting the idea of a credible interval for a discrete parameter. We firstly illustrate our definition with an example, and then formalise it below. Suppose that given data y_0 simulated from model $m_0 \in \{1, 2, 3\}$, estimated posterior probabilities are 0.7, 0.2 and 0.1. This could be viewed as defining three credible intervals; a 70% credible interval that $m = 1$ etc. We would like to investigate coverage in the following sense: given a 70% interval for $m = 1$ produced by some (m_0, y_0) pair, there is a 70% probability of it containing m_0 . A technical difficulty is that the probability of a pair producing a 70% credible interval is typically zero. This difficulty can be avoided by requiring the following condition to hold for every $a < b$ e.g. $a = 0.69$ and $b = 0.71$: Consider all y_0 such that the estimated probability of $m = 1$ is between a and b . Conditioning on this, the probability of $m_0 = 1$ also lies between a and b .

Definition of coverage property: Let $g(m|y)$ be a mass function approximating the posterior and $G_y(m)$ the corresponding distribution function. Given $I = [a, b] \subseteq [0, 1]$, define $A(m, I) = \{y | g(m|y) \in I\}$. We say g satisfies the coverage property with respect to distribution $H(m_0, y_0)$ if, for all $i \in \{1, 2, \dots, M\}$ and I , either

$$\Pr(y_0 \in A(i, I)) = 0, \quad \text{or} \quad (3)$$

$$\Pr(m_0 = i | y_0 \in A(i, I)) \in I. \quad (4)$$

Similar arguments to the parameter inference case show that the posterior satisfies coverage when $h(m_0|y_0) = \pi(m_0|y_0)$, but the prior satisfies coverage when $h(m_0) = p(m_0)$.

Result 4 *The posterior, $p(m|y)$, satisfies coverage with respect to any distribution $H(m_0, y_0)$*

with conditional mass function $h(m_0|y_0) = p(m_0|y_0)$.

Proof in Appendix.

Result 5 *The prior, $p(m)$, satisfies coverage with respect to any distribution $H(m_0, y_0)$ with marginal mass function $h(m_0) = p(m_0)$.*

Proof in Appendix.

This means that the natural choice of $h(m_0, y_0) = p(m_0, y_0)$ (where $p(m_0, y_0) = p(y_0|m_0)p_0(m_0) = p(m_0|y_0)p(y_0)$) is not suitable. As before, our proposed solution is to truncate this distribution on (m_0, y_0) , so that $h(m_0, y_0) = p(m_0, y_0)\mathbb{I}[y_0 \in A]$.

As before, the above definition of coverage for model inference is not directly testable. Below we give an equivalent (under weak technical conditions) form which is.

Result 6 *Define $z_0^{(i)} = g(i|y_0)$. Assume that for all $i \in \{1, 2, \dots, M\}$, the measure $\mathcal{Z}^{(i)}$ on $z_0^{(i)}$ induced by the distribution $H(m_0, y_0)$ is not a singular continuous distribution with respect to Lebesgue measure. Coverage then holds with respect to $H(m_0, y_0)$ if and only if, for all i*

$$\Pr(m_0 = i | z_0^{(i)} = w) = w \quad (5)$$

holds for almost all w with respect to $\mathcal{Z}^{(i)}$.

Proof in Appendix.

3 Method

In this Section we discuss how to construct diagnostics based on the coverage property. In principle, this is simply a matter of repeatedly constructing an ABC posterior approximation, $\pi_{\text{ABC}}(m, \theta|y_0)$, for known values of $(m_0, \theta_0, y_0) \sim H(m_0, \theta_0, y_0)$, computing p -values and estimated model probabilities, and then testing whether the conditions (2) and (5) hold. This can be repeated for many ϵ values until a suitable choice is found. However, simulating datasets for a single ABC analysis is typically computationally expensive. As such, we reuse the same simulations for each ABC analysis, along the lines of Algorithm 1, as is common for ABC diagnostics (e.g. Blum et al. (2013)). Simulations in Section 4 indicate this makes little difference to the results.

In the following, we first present the full algorithm, including how to generate $(m_0, \theta_0, y_0) \sim H(m_0, \theta_0, y_0)$. We then describe several test statistics and diagnostic plots to allow an assessment of whether the conditions (2) and (5) hold. As in the previous Section, the details are presented in terms of y rather than s for notational simplicity. R code to implement these methods has been made available as part of the `abc` package (Csilléry et al., 2012).

3.1 Algorithm

ABC coverage diagnostics

1. Determine integers $N > 0$, $c > 0$ and candidate values of ϵ : $\epsilon_1 > \epsilon_2 > \dots \epsilon_q \geq 0$.
 2. Simulate a set $\mathbb{U} = \{(m_i, \theta_i, y_i) | i = 1, \dots, N\}$ of independent realisations of (m, θ, y) from $\pi(y|\theta, m)\pi(\theta|m)p(m)$.
 3. Select $\mathbb{V} \subseteq \mathbb{U}$ containing the c realisations that minimise $\|y_i - y_{\text{obs}}\|$.
 4. For each $(m_0, \theta_0, y_0) \in \mathbb{V}$ and for $j = 1, \dots, q$:
 - (a) Let $\mathbb{W} = \mathbb{U} \setminus (m_0, \theta_0, y_0)$.
 - (b) Find the subset of $\mathbb{W}$ such that $\|y_i - y_{\text{obs}}\| \leq \epsilon_j$.
 - (c) (Optional) Perform regression-adjustment post-processing.
 - (d) Record p -values and estimated model probabilities.
 5. Construct plots of diagnostic statistics versus ϵ .
-

Algorithm 2: An algorithm to diagnose coverage for ABC as a function of kernel scale parameter $\epsilon > 0$. In the case of a single model, modify $(m, \theta, y) \rightarrow (\theta, y)$ and $\pi(y|\theta, m)\pi(\theta|m)p(m) \rightarrow \pi(y|\theta)\pi(\theta)$ in the obvious way.

The algorithm for diagnosing coverage of the ABC approximation $\pi_{ABC}(m, \theta|y_{\text{obs}})$ (or $\pi_{ABC}(\theta|y_{\text{obs}})$) is presented in Algorithm 2. The set $\mathbb{V}$ is a sample of size c from the distribution $h(m_0, \theta_0, y_0) = \pi(y_0|\theta_0, m_0)\pi(\theta_0|m_0)p(m_0)\mathbb{I}[y_0 \in A]$, where $A = \{y : \|y - y_{\text{obs}}\| \leq \delta\}$ for some δ determined by c . Each element of $\mathbb{V}$ is taken as the known values (m_0, θ_0, y_0) in turn, and the ABC posterior approximation estimated for a range of kernel scale parameters, ϵ .

Increasing c , the number of known values of (m_0, θ_i, y_0) , will improve the power of the tests of coverage. However the tradeoff is a greater computing cost, and that A becomes less concentrated around y_{obs} , so the risk of the prior satisfying the coverage property increases.

The final choice is left to the user. However, we note that c can be increased (or decreased) based on preliminary findings. We investigate various values of c by simulation in Section 4, and based on this suggest $c = 200$ as a default.

3.2 P -values and model probabilities

For scalar θ , we require a p -value estimate of (2) under Result 3, based on a posterior sample $\theta^{(1)}, \theta^{(2)}, \dots, \theta^{(n)}$. Here, we use $p = (1 + \sum_{j=1}^n \mathbb{I}(\theta^{(j)} < \theta_0)) / (2 + n)$, which is equivalent to the posterior mean for the binomial probability that $\theta_j < \theta_0$ under a uniform prior. This choice eliminates the occurrence of extreme values ($p = 0$ or 1), which can overly influence some test statistics. For multivariate θ , we record a p -value estimate for each parameter.

Given a posterior sample of model indicators $m^{(1)}, m^{(2)}, \dots, m^{(n)}$, a straightforward estimate of the posterior probability of model i is the proportion which equal this: $g(m = i|y_0) = \sum_{j=1}^n \mathbb{I}(m^{(j)} = i) / n$. Alternatively, regression-adjusted post-processing produces estimated posterior model probabilities directly (Beaumont, 2008).

Removing an element of $\mathbb{U}$ in step 4a of Algorithm 2 can slightly bias the estimated ABC model probabilities. For example, let $d = \sum_{i=1}^n \mathbb{I}[m_i = 1]$. For $\epsilon = \infty$, $g(m = 1|y_0) = (d - 1) / (n - 1)$ for $m_0 = 1$ and $d / (n - 1)$ otherwise. This dependence of $g(\cdot|y_0)$ on m_0 causes unwanted behaviour in some diagnostics. To mitigate this, we reweight the model probability estimates to use the empirical prior model weights from $\mathbb{U}$ rather than $\mathbb{W}$. That is, given estimated model probabilities $g(m_i|y_0)$ we adjust these to $\tilde{g}(m = i|y_0) \propto g(m = i|y_0) h_i(\mathbb{U}) / h_i(\mathbb{W})$, where $h_i(\cdot)$ gives the proportion of realisations from model i in the supplied set. No similar correction of parameter estimates was found to be necessary.

3.3 Diagnostic statistics

For each parameter and value of ϵ we will have c replicated p -values, $p_1, p_2, \dots, p_c$. We treat these as independent, although there may be mild dependence induced by Algorithm 2. Under the coverage property these will be distributed as $U(0, 1)$ (Result 3). There are a

number of tests for uniformity. Cook et al. (2006) used the diagnostic statistic

$$X^2 = \sum_{i=1}^c (\Phi^{-1}(p_i))^2, \quad (6)$$

where Φ is the standard normal distribution function. When the p_i values are independent $U(0, 1)$ draws, $X^2 \sim \chi_c^2$, which allows the calculation of a p -value for coverage (we report the p -value for a two-tailed test). We note that this statistic is unchanged if some p_i values are replaced with $1 - p_i$; it does not test for symmetry of the distribution around 0.5. This can cause problems in practice. An example based on real data is the top left histogram of Figure S5 which displays p_i values that are clearly not uniform but receive a p -value of 0.75 from this diagnostic statistic.

An alternative used by Wegmann et al. (2009, 2010) is the Kolmogorov-Smirnoff test statistic

$$Y = \sup_x |F_c(x) - F(x)|, \quad (7)$$

where $F_c(x)$ is the empirical distribution function of $p_1, p_2, \dots, p_c$ and $F(x)$ is the $U(0, 1)$ distribution function. The distribution of Y if $p_i \sim U(0, 1)$ is known in exact and asymptotic forms (Durbin, 1973) and can be used to calculate a p -value for coverage (using a one-tailed test). Our p_i values are not drawn from continuous distributions, but rather discrete distributions based on the number of posterior samples. However asymptotic p -values based on the continuous distribution can still be calculated and will be of the correct order of magnitude, which suffices for their purpose as a rough diagnostic guide. If more accurate p -values are required, Monte Carlo estimation is possible but more time consuming.

For model inference diagnostics, we focus on the binary case of model $m = i$ and model $m \neq i$, for each $i \in \{1, \dots, M\}$. For each ϵ , Algorithm 2 is run on a sequence of y_0 values, $y_{0,1}, y_{0,2}, \dots, y_{0,c}$, generated from m_0 values, $m_{0,1}, m_{0,2}, \dots, m_{0,c}$, to produce z values, $z_1, z_2, \dots, z_c$, where z_j is the estimated probability of model i : $\Pr_{\text{ABC}}(m = i | y_{0,j})$. Define $q_j = \mathbb{I}(m_{0,j} = i)$. Following Result 6, we wish to test the coverage hypothesis that $q_j \sim$

Bernoulli(z_j), where all q_j values are assumed independent, as before.

A simple diagnostic statistic is the proportion of times model i occurs,

$$U = c^{-1} \sum_{j=1}^c q_j.$$

A central limit theorem holds for the distribution of U under the null hypothesis of coverage, conditional on the z_j values. However, this can be a poor approximation when some z_j values are close to 0 or 1. Instead, we construct the null distribution by Monte Carlo methods to estimate the p -value for coverage (using a two-tailed test). To improve the stability of the resulting p -values, we use the same random seed across different ϵ values.

A drawback of U is that highly unlikely q_j values, such as $q_j = 1$ when $z_j = 10^{-6}$ provide strong evidence against coverage, but do not contribute more to U . As an alternative, we can consider the log-likelihood,

$$V = \sum_{j=1}^c [q_j \log z_j + (1 - q_j) \log(1 - z_j)], \quad (8)$$

with p -values for coverage (using a two-tailed test) again calculated by Monte Carlo simulation. A drawback of this statistic is that V is constant regardless of the q_i values if $z_j \equiv 0.5$, and so coverage cannot be rejected. Also note a similar statistic is to use the log-likelihood of c independent discrete random variables, $W = \sum_{j=1}^c \log \text{Pr}_{\text{ABC}}(m = m_{0,j} | y_{0,j})$. This tests coverage of all models.

A problem with these statistics is that they can be insensitive to departures from coverage which vary in nature with q_j . It is difficult to define a statistic which is flexible enough to detect such problems for all possible (q_j, z_j) sequences. Seillier-Moiseiwitsch and Dawid (1993) present a portmanteau statistic combining tests on a partition chosen for a specific dataset, but in our experience such a statistic is hard to adapt to work generally. As such, we advise that checking diagnostic plots is particularly important. Should these show poor performance of general purpose diagnostic statistics, they may motivate a better choice

specific to the problem of interest.

3.4 Diagnostic plots

For parameter inference, many standard diagnostic plots can be used to assess uniformity of $p_1, p_2, \dots, p_c$, such as histograms and probability plots.

For model inference, we present a diagnostic plot, an example of which is shown in Figure 7. Based on an equally-spaced partition of $[0, 1]$ into subintervals, S , we estimate $\Pr(q_j = 1 | z_j \in S)$ for each subinterval, by Bayesian inference for a binomial rate using a uniform prior. The diagnostic plot displays each posterior mean and 95% credible interval. Under coverage, each credible interval should include some of the associated z_j interval with high probability. The plot illustrates whether the coverage property appears to hold for each interval individually. This approach is similar to the Seillier-Moiseiwitsch and Dawid (1993) portmanteau statistic described above, but without the need to combine the results into a single statistic. Indeed, they also propose a “coverage plot,” similarly plotting point estimates for several partitions.

4 Analyses

4.1 Simulated example

We now examine how our coverage diagnostics perform in a simple simulated example. We consider that 100 data points are drawn independently from either a $N(0, 1)$ or a $gk(0, 1, g, 0)$ model, which are equally likely a priori. (For details on the g -and- k distribution see e.g. Drovandi and Pettitt 2011.) Inference can be split into binary model choice, and inference for the unknown parameter, $g > 0$, which has a $U(0, 4)$ prior. Observed data, y_{obs} , was drawn from the g -and- k model with $g = 0.2$. We base our ABC analysis on the median and upper and lower quartiles as summary statistics, so we are interested in determining how well the ABC approximation $\pi_{\text{ABC}}(m, \theta | s_{\text{obs}})$ represents the posterior $\pi(m, \theta | s)$. For the

analyses in this paper, we use weighted Euclidean distance $\|a - b\| = [\sum_j (a_j - b_j)^2 / v_j^2]^{1/2}$, where v_j^2 is the prior predictive variance of the j -th summary statistic, estimated from the set $\mathbb{W}$ in Algorithm 2.

For analysis, we construct the set $\mathbb{U}$ from 2×10^6 simulated (m, θ, y) triples, half from each model. Figure 1 shows coverage diagnostic p -values from the statistics U , V , X^2 and Y as a function of ϵ , and for parameter (left panels) and model (right panels) inference respectively. For parameter inference, we only present results from the g -and- k model. The top panels show diagnostics when the set of known values, $\mathbb{V} = \{(m_0, \theta_0, s_0)\}$, is a random sample of size $c = 200$ from $\mathbb{U}$ (i.e. the prior), and the middle and bottom panels when $\mathbb{V}$ consists of the $c = 200$ samples with s closest to s_{obs} .

The top panels in Figure 1 support Results 2 and 5, in that when $(m_0, \theta_0, s_0) \sim \pi(s|\theta, m)\pi(\theta|m)p(m)$ are drawn from the prior, then coverage holds both for the prior (i.e. large ϵ values) and the posterior (i.e. small ϵ values). When $(m_0, \theta_0, s_0) \sim \pi(s|\theta, m)\pi(\theta|m)p(m)\mathbb{I}(\|s_0 - s_{\text{obs}}\| \leq \delta)$ are drawn from the truncated prior (Figure 1, middle panels), then coverage does not hold for the prior. Note that the statistic U does not detect any deviation from coverage here, as discussed in Section 3.3. Figure 1 also illustrates our earlier point, particularly in the case of parameter inference, that requiring coverage to hold for the prior is more demanding than requiring coverage to hold for the truncated prior. This is evidenced by the upturn in p -values only occurring at lower ϵ values when $\mathbb{V}$ is drawn from the prior, compared to the relatively larger values of ϵ when $\mathbb{V}$ is drawn from the truncated prior.

For comparison, the procedure with $\mathbb{V}$ drawn from the truncated prior was repeated by resimulating $\mathbb{W}$ for each ABC analysis, thereby removing any effects of reusing (m, θ, s) samples, albeit at far greater computational expense. Figure 1 (bottom panels) shows that the results are nearly identical to those obtained using Algorithm 2. Further, we repeated our analysis with $c = 100$ and $c = 500$, and obtained qualitatively similar results (see Figures S1 and S2 in the Supplementary Information), suggesting that the choice of c is not crucial to drawing the correct inferences.

Figures 2 and 3 show diagnostic plots to investigate coverage in more detail for $\epsilon = 0.28, 1.5, 13$. These again demonstrate that coverage approximately holds for large ϵ when $\mathbb{V}$ is drawn from the prior, but not from the truncated prior. They also provide insight into disagreements between the statistics within some panels in Figure 1. For parameter inference, in the top left panel of Figure 1, there is less than clear agreement about whether coverage roughly holds for the smallest ϵ values. The top left panel of Figure 2 confirms that the p -value histogram has a non-uniform shape in this case. For model inference, the top right panel of Figure 1 suggests no deviation from coverage for any ϵ for the U statistic. However, the top centre panel of Figure 3 illustrates that this is not correct.

Our interpretation of these results is that $\epsilon \leq 0.28$ is sufficient to achieve approximate coverage for both parameter inference and model selection in the case where $\mathbb{V}$ is drawn from the truncated prior. Coverage does not hold (for small ϵ) when $\mathbb{V}$ is drawn from the prior, which represents a stricter condition. However, the former case relates more to the dataset and analysis of interest.

4.2 Application in human demographic history

Sjödín et al. (2012) detail an ABC analysis of genetic data to choose between three demographic models of human history: null, bottleneck and fragmentation models. Each model contains 9 unknown parameters: b , d , n , N_0 , N_A , N_B , T_b , T_g and T_{dur} . Their analysis used 10^5 simulations from each model, and they accepted the 0.5% of simulations minimising $\|s - s_{\text{obs}}\|$, corresponding to $\epsilon = 1.36$. Sjödín et al. (2012) then used regression post-processing for parameter and model inference. We use the same experimental setup to evaluate the implemented value of ϵ , and also to determine whether regression-adjusted post-processing improved the results.

Figure 4 shows parameter inference diagnostics for d , N_0 , N_A and T_b , without regression post-processing (see Supporting Information Figure S3 for the same plot for the remaining parameters). There is occasional significant disagreement between the statistics. However,

overall it is apparent that coverage is not attained for any ϵ , apart from perhaps T_{dur} . P -value histograms for $\epsilon = 1.36$ (Figure S5) confirm that in most cases there is clear deviation from coverage. For model choice, both statistics agree that coverage is not attained (Figure 6; left panels) and diagnostic plots for $\epsilon = 1.36$ confirm this (Figure 7; left panels).

Regression post-processing was performed by conditional heteroskedastic, local-linear regression (Blum and François, 2010) for parameter inference, and multinomial logistic regression (Beaumont, 2008) for model inference. The regression post-processing greatly improves the results. The model inference statistics (Figure 6; right panels) now suggest that coverage holds for any choice of ϵ investigated. The same is true of many parameters, although for others, coverage appears to hold only for smaller ϵ (Figure 5, Figure S4). Diagnostic plots for parameter and model inference (Figure S6 and 7; right panels) produced for $\epsilon = 1.36$ suggest that coverage is approximately achieved, except for some small concerns remaining for some parameters (e.g. N_B and T_{dur}).

On the whole, and with only a few minor caveats, our investigation largely validates the choice of ϵ , and the use of regression post-processing by Sjödin et al. (2012).

5 Discussion

We have presented a method for validating whether an ABC analysis contains significant approximation error based on assessment of the coverage property. The method can be used to determine the kernel scale parameter, ϵ , via simple diagnostic plots. We have used this method in a re-analysis of human demographic data (Sjödin et al., 2012), validating the choice of ϵ and the use of regression-adjustment post-processing in that study.

Our methodology draws on several previous approaches. In particular, Wegmann et al. (2009, 2010) use a similar scheme for validating ABC parameter inference, and suggest using the Y diagnostic statistic (7). Also, Cook et al. (2006) employ a similar idea for Bayesian software testing using the X^2 statistic (6). Our contribution here is to provide: results on

the choice of (m_0, θ_0, y_0) samples to use, a description of a general purpose methodology (and R code to implement it), and evidence that many diagnostic statistics are not fully trustworthy and should be supplemented with diagnostic plots. We also extend the coverage property definition and the validation methodology to cover model inference, incorporating ideas from Seillier-Moiseiwitsch and Dawid (1993).

Our approach only aims to determine whether $\pi_{\text{ABC}}(m, \theta | y_{\text{obs}})$ is a good approximation of $\pi(m, \theta | y_{\text{obs}})$, or whether $\pi_{\text{ABC}}(m, \theta | s_{\text{obs}})$ is a good approximation of $\pi(m, \theta | s_{\text{obs}})$. In order to use the coverage property to assess the approximation impact of summary statistics, in addition to that of ϵ , it would be necessary to choose the set $\mathbb{V}$ to consist of those (m, θ, y) that minimise $\|y - y_{\text{obs}}\|$, and then perform the rest of the analysis based on using $(m, \theta, S(y))$ rather than (m, θ, y) . Further investigation would be required to see if this is a practical approach.

One word of caution: when a summary statistic is not informative for a parameter, so that $\pi_{\text{ABC}}(\theta | s) = \pi(\theta | s) = \pi(\theta)$, then our diagnostics will support a good posterior approximation for any value of ϵ . This is, of course, the correct result, however it should not be misconstrued as any support of information content in s for θ . Additionally, the above diagnostics are evaluated for each parameter separately within a multivariate parameter θ . Hence, there is the possibility of diagnosing a good posterior approximation for all posterior margins, but not for the joint distribution of model parameters within any model. This could be resolved by constructing a suitable multivariate diagnostic and test.

R code for implementing Algorithm 2 can be found in the **abc** package, which is freely available on the CRAN.

Preprint note: The R code is currently being incorporated into the above package. In the meantime, it is directly available from <http://www.maths.lancs.ac.uk/~prangle/pub.html>

Appendix: Proofs

Here we provide proofs of Results 1–6 presented in Section 2. Denote by $F(m, \theta, y)$ the joint distribution function defined by $\pi(y|\theta, m)\pi(\theta|m)p(m)$. We will also use F to denote the associated marginal and conditional distributions.

Proof of Result 1: We have $g(\theta|y) = \pi(\theta|y)$ and $h(\theta_0|y_0) = \pi(\theta_0|y_0)$. In this case, $G_y(\theta) = F(\theta|y) = H(\theta|y)$. Hence

$$\begin{aligned} \Pr(\theta_0 \in C(y_0, \alpha)|y_0) &= \Pr(G_{y_0}(\theta_0) \in B(\alpha)|y_0) = \alpha \\ \Rightarrow \Pr(\theta_0 \in C(y_0, \alpha)) &= \mathbb{E}_{G(y_0)}[\Pr(\theta_0 \in C(y_0, \alpha)|y_0)] = \alpha. \end{aligned}$$

Proof of Result 2: We have $g(\theta|y) = \pi(\theta)$ and $h(\theta_0) = \pi(\theta_0)$. In this case, $G_y(\theta) = F(\theta) = H(\theta)$. Hence

$$\Pr(\theta_0 \in C(y_0, \alpha)) = \Pr(G_{y_0}(\theta_0) \in B(\alpha)) = \alpha.$$

Proof of Result 3: First assume that coverage holds. Let $B(\alpha) = [0, \alpha]$. Then

$$\alpha = \Pr(\theta_0 \in C(y_0, \alpha)) = \Pr(p_0 \in [0, \alpha]), \quad (\text{applying } G_{y_0} \text{ to the event})$$

so the distribution function of p_0 equals that of a $U(0, 1)$ distribution. For the converse, now assume that $p_0 \sim U(0, 1)$. Then

$$\alpha = \Pr(p_0 \in B(\alpha)) = \Pr(\theta_0 \in C(y_0, \alpha)), \quad (\text{applying } G_{y_0}^{-1} \text{ to the event})$$

which is the condition needed for coverage.

Proof of Result 4: We have $g(m|y) = p(m|y)$ and $h(m_0|y_0) = p(m_0|y_0)$. In this case, $g(m|y) = p(m|y) = h(m|y)$. Fix some i and I such that (3) does not hold, and write A for

$A(i, I)$. Then for $y_0 \in A$, $h(i|y_0) = g(i|y_0) \in I$. Thus $\Pr(m_0 = i|y_0 \in A) = \mathbb{E}_{H(y_0)}[h(i|y_0)] \in I$, where $H(y_0)$ denotes the marginal distribution of H . Hence (4) holds.

Proof of Result 5: We have $g(m|y) = p(m)$ and $h(m_0) = p(m_0)$. In this case, $g(m|y) = p(m) = h(m)$. Fix some i and I and write A for $A(i, I)$. Consider first that $p(i) \notin I$. Then $A = \emptyset$ and so (3) holds. Suppose instead that $p(i) \in I$. Then A is the set of all possible y values, and $\Pr(m_0 = i|y_0 \in A) = \Pr(m_0 = i) = p(i) \in I$. Hence (4) holds.

Proof of Result 6: Assume that, for all i , (5) holds for almost all w (with respect to $\mathcal{Z}^{(i)}$ as defined in the statement of this result). Fix some i and I , and consider the case where (3) does not hold. Then

$$\begin{aligned} \Pr(m_0 = i|y_0 \in A(i, I)) &= \Pr(m_0 = i|z_0 \in I) \\ &= \mathbb{E}_{Z'(z_0)}[p(i|z_0)] \\ &= \mathbb{E}_{Z'(z_0)}[z_0] \in I, \end{aligned}$$

where $Z'(z_0)$ is the marginal distribution of z_0 truncated to I . Thus, coverage holds.

Next assume coverage with respect to H , and fix some i . For any w such that $\Pr(z_0 = w) > 0$, it is immediate that (5) follows. Define $I_\varepsilon(w) = [w - \varepsilon, w + \varepsilon] \cap [0, 1]$. It suffices to prove that (5) holds for w such that $\Pr(z_0 \in I_\varepsilon(w)) > 0$ for all $\varepsilon > 0$. (The set of other w values has probability zero, as each w lies within an interval of zero probability.) Fix such a w with $\Pr(z_0 = w) = 0$. Note that $z_0 \in I_\varepsilon(w)$ represents the same event as $y_0 \in A_\varepsilon := A(i, I_\varepsilon(w))$. From the assumption on w , (3) is false for $I = I_\varepsilon(w)$ and $\varepsilon > 0$. Hence (4) must hold i.e.

$$\Pr(m_0 = i|y_0 \in A_\varepsilon) \in I_\varepsilon(w) \quad \text{for } \varepsilon > 0.$$

The left hand side of this equals

$$\Pr(m_0 = i | z_0 \in I_\varepsilon(w)) = E_{Z'_\varepsilon(z_0)}[\Pr(m_0 = i | z_0)],$$

where $Z'_\varepsilon(z_0)$ is the marginal distribution of z_0 truncated to $I_\varepsilon(w)$. Thus

$$E_{Z'_\varepsilon(z_0)}[\Pr(m_0 = i | z_0)] \in I_\varepsilon(w).$$

It follows by the Lebesgue differentiation theorem, using the assumption on z_0 assumed in the statement of Result 6, that $\Pr(m_0 = i | z_0 = w) = w$ for almost all w , as required.

References

- Aeschbacher, S., Beaumont, M. A., and Futschik, A. (2012). A novel approach for choosing summary statistics in approximate Bayesian computation. *Genetics*, 192:1027–1047.
- Beaumont, M. A. (2008). Joint determination of topology, divergence time, and immigration in population trees. In *Renfrew C Matsumura S, Forster P, editor, Simulation, Genetics and Human Prehistory, McDonald Institute Monographs*, pages 134–154. McDonald Institute Monographs.
- Beaumont, M. A. (2010). Approximate Bayesian computation in evolution and ecology. *Annual Review of Ecology, Evolution, and Systematics*, 41(1).
- Beaumont, M. A., Zhang, W., and Balding, D. J. (2002). Approximate Bayesian computation in population genetics. *Genetics*, 162:2025–2035.
- Bertorelle, G., Benazzo, A., and Mona, S. (2010). ABC as a flexible framework to estimate demography over space and time: some cons, many pros. *Molecular Ecology*, 19(13):2609–2625.

- Blum, M. G. B. (2010a). Approximate Bayesian computation: A nonparametric perspective. *Journal of the American Statistical Association*, 105:1178–1187.
- Blum, M. G. B. (2010b). Choosing the summary statistics and the acceptance rate in approximate Bayesian computation. In *Proceedings of COMPSTAT'2010*, pages 47–56.
- Blum, M. G. B. and François, O. (2010). Non-linear regression models for approximate Bayesian computation. *Statistics and Computing*, 20:63–73.
- Blum, M. G. B., Nunes, M., Prangle, D., and Sisson, S. A. (2013). A comparative review of dimension reduction methods in approximate Bayesian computation. *Statistical Science (forthcoming)*.
- Bortot, P., Coles, S. G., and Sisson, S. A. (2007). Inference for stereological extremes. *Journal of the American Statistical Association*, 102:84–92.
- Cook, S. R., Gelman, A., and Rubin, D. B. (2006). Validation of software for Bayesian models using posterior quantiles. *J. Comp. Grap. Stat.*, 15(3):675–692.
- Csilléry, K., Blum, M. G. B., Gaggiotti, O. E., and François, O. (2010). Approximate Bayesian computation (ABC) in practice. *Trends in Ecology & Evolution*, 25(7):410–418.
- Csilléry, K., François, O., and Blum, M. G. B. (2012). abc: An R package for approximate Bayesian computation (ABC). *Methods in Ecology and Evolution*, 3(3):475–479.
- Dawid, A. P. (1984). Statistical theory: The prequential approach. *Journal of the Royal Statistical Society: Series A*, 147:278–292.
- Drovandi, C. C. and Pettitt, A. N. (2011). Likelihood-free Bayesian estimation of multivariate quantile distributions. *Computational Statistics & Data Analysis*, 55:2541–2556.
- Durbin, J. (1973). *Distribution theory for tests based on the sample distribution function*. Society for Industrial Mathematics.

- Faisai, M., Futschick, A., and Hussain, I. (2013). A new approach to choose acceptance cutoff for approximate Bayesian computation. *Journal of Applied Statistics*, page in press.
- Gneiting, T., Balabdaoui, F., and Raftery, A. E. (2007). Probabilistic forecasts, calibration and sharpness. *Journal of the Royal Statistical Society: Series B*, 69:243–268.
- Grelaud, A., Robert, C., Marin, J.-M., Rodolphe, F., and Taly, J. F. (2009). ABC likelihood-free methods for model choice in Gibbs random fields. *Bayesian Analysis*, 4(2):317–336.
- Marin, J.-M., Pudlo, P., Robert, C. P., and Ryder, R. J. (2012). Approximate Bayesian computational methods. *Statistics and Computing*, 22:1167–1180.
- Menéndez, P., Fan, Y., Garthwaite, P. H., and Sisson, S. A. (2012). Simultaneous adjustment of bias and coverage probabilities for confidence intervals. <http://arxiv.org/abs/1210.3405>.
- Ratmann, O., Andrieu, C., Wiuf, C., and Richardson, S. (2009). Model criticism based on likelihood-free inference, with an application to protein network evolution. *Proceedings of the National Academy of Sciences*, 106(26):10576.
- Seillier-Moiseiwitsch, F. and Dawid, A. P. (1993). On testing the validity of sequential probability forecasts. *Journal of the American Statistical Association*, 88:355–359.
- Sisson, S. A. and Fan, Y. (2011). Likelihood-free Markov chain Monte Carlo. In Brooks, S. P., Gelman, A., Jones, G., and Meng, X.-L., editors, *Handbook of Markov Chain Monte Carlo*, pages 319–341. Chapman and Hall/CRC Press.
- Sjödin, P., Sjöstrand, A. E., Jakobsson, M., and Blum, M. G. B. (2012). Resequencing data provide no evidence for a human bottleneck in africa during the penultimate glacial period. *Molecular Biology and Evolution*, 29:1851–1860.
- Wegmann, D., Leuenberger, C., and Excoffier, L. (2009). Efficient approximate Bayesian computation coupled with Markov chain Monte Carlo without likelihood. *Genetics*, 182:1207–1218.

Wegmann, D., Leuenberger, C., Neuenschwander, S., and Excoffier, L. (2010). ABCtoolbox: a versatile toolkit for approximate Bayesian computations. *BMC bioinformatics*, 11(1):116.




Figure 1: Plots of ϵ against coverage p -values for the $N(0, 1)$ / g -and- k example. Left panels correspond to parameter inference, and right panels for model inference. Top panels indicate when V consists of $c = 200$ samples drawn from the prior; middle panels when V is constructed from the $c = 200$ samples in which s is closest to s_{obs} . Top and middle panels use Algorithm 2 which re-uses (m, θ, s) samples throughout the analyses. Bottom panels simulate new (m, θ, s) samples for each ABC analysis.




Figure 2: Histograms of the $c = 200$ p_0 values for the parameter g in the $N(0, 1)$ / g -and- k example, for $\epsilon = 0.28, 1.5, 13$. In the top panels $\mathbb{V}$ is drawn from the prior; in the bottom panels $\mathbb{V}$ is drawn from the truncated prior. Columns indicate different ϵ values.




Figure 3: Model inference diagnostics in the $N(0, 1)$ / g -and- k example, for $\epsilon = 0.28, 1.5, 13$. In the top panels $\mathbb{V}$ is drawn from the prior; in the bottom panels $\mathbb{V}$ is drawn from the truncated prior. Columns indicate different ϵ values. Each panel shows the observed and predicted (under coverage) model probabilities for the $N(0, 1)$ model, including a 95% credible interval for the predictions.




Figure 4: Plots of ϵ against coverage p -values for the parameters d , N_0 , N_A and T_b in the human demographic history analysis. Regression-adjusted post-processing is not implemented. Rows correspond to individual parameters; columns correspond to the three models.




Figure 5: Plots of ϵ against coverage p -values for the parameters d , N_0 , N_A and T_b in the human demographic history analysis. Regression-adjusted post-processing has been implemented. Rows correspond to individual parameters; columns correspond to the three models.




Figure 6: Plots of ϵ against coverage p -values in the human demographic history analysis. Rows correspond to the three models; columns correspond to the implementation of regression-adjustment post-processing.




Figure 7: Model inference diagnostics in the human demographic history analysis. Rows represent (top) the null, (middle) bottleneck and (bottom) fragmentation models; columns correspond to the implementation of regression-adjustment post-processing. Each panel shows the observed and predicted (under coverage) model probabilities for each model, including a 95% credible interval for the predictions.




Figure 8: (Supplementary Figure 1): As Figure 1 (main text), but with $c = 500$.




Figure 9: (Supplementary Figure 2): As Figure 1 (main text), but with $c = 100$.




Figure 10: (Supplementary Figure 3): As for Figure 4 (main text), but for the remaining model parameters T_g , b , N_B , T_{dur} and n .




Figure 11: (Supplementary Figure 4): As for Figure 5 (main text), but for the remaining model parameters T_g , b , N_B , T_{dur} and n .




Figure 12: (Supplementary Figure 5): Histograms of the $c = 200 p_0$ values for the parameters d , N_0 , N_A and T_b in the human demographic history analysis, with $\epsilon = 1.36$. Regression-adjusted post-processing is not implemented. Rows correspond to individual parameters; columns correspond to the three models.




Figure 13: (Supplementary Figure 6): Histograms of the $c = 200$ p_0 values for the parameters d , N_0 , N_A and T_b in the human demographic history analysis, with $\epsilon = 1.36$. Regression-adjusted post-processing has been implemented. Rows correspond to individual parameters; columns correspond to the three models.